

Consensus on current management of endometriosis

Neil P. Johnson^{1,2,3,*} and Lone Hummelshoj¹, for the World Endometriosis Society Montpellier Consortium[†]

¹World Endometriosis Society, 89 Southgate Road, London N1 3JS, UK ²Repromed Auckland, Auckland, New Zealand ³University of Auckland, Auckland, New Zealand

*Correspondence address. Tel: +44-77-1006-5164; E-mail: wes@endometriosis.ca

Submitted on December 16, 2012; resubmitted on January 25, 2013; accepted on February 8, 2013

STUDY QUESTION: Is there a global consensus on the management of endometriosis that considers the views of women with endometriosis?

SUMMARY ANSWER: It was possible to produce an international consensus statement on the current management of endometriosis through engagement of representatives of national and international, medical and non-medical societies with an interest in endometriosis.

WHAT IS KNOWN ALREADY: Management of endometriosis anywhere in the world has been based partially on evidence-based practices and partially on unsubstantiated therapies and approaches. Several guidelines have been developed by a number of national and international bodies, yet areas of controversy and uncertainty remain, not least due to a paucity of firm evidence.

STUDY DESIGN, SIZE, DURATION: A consensus meeting, in conjunction with a pre- and post-meeting process, was undertaken.

PARTICIPANTS/MATERIALS, SETTING, METHODS: A consensus meeting was held on 8 September 2011, in conjunction with the 11th World Congress on Endometriosis in Montpellier, France. A rigorous pre- and post-meeting process, involving 56 representatives of 34 national and international, medical and non-medical organizations from a range of disciplines, led to this consensus statement.

MAIN RESULTS AND THE ROLE OF CHANCE: A total of 69 consensus statements were developed. Seven statements had unanimous consensus; however, none of the statements were made without expression of a caveat about the strength of the statement or the statement itself. Only two statements failed to achieve majority consensus. The statements covered global considerations, the role of endometriosis organizations, support groups, centres or networks of expertise, the impact of endometriosis throughout a woman's life course, and a full range of treatment options for pain, infertility and other symptoms related to endometriosis.

LIMITATIONS, REASONS FOR CAUTION: This consensus process differed from that of formal guideline development. A different group of international experts from those participating in this process would likely have yielded subtly different consensus statements.

WIDER IMPLICATIONS OF THE FINDINGS: This is the first time that a large, global, consortium, representing 34 major stakeholding organizations from five continents, has convened to systematically evaluate the best available current evidence on the management of endometriosis, and to reach consensus. In addition to 18 international medical organizations, representatives from 16 national endometriosis organizations were involved, including lay support groups, thus generating input from women who suffer from endometriosis.

STUDY FUNDING/COMPETING INTEREST(S): The World Endometriosis Society commissioned and hosted the consensus meeting. Financial support for participants to attend the meeting was provided by the organizations that they represented. There was no other specific funding for this consensus process. Full disclosures of all participants are presented herein.

Key words: endometriosis / evidence based / management / WES Montpellier Consortium / World Endometriosis Society

[†] The complete list of people representing The World Endometriosis Society Montpellier Consortium is given in Appendix.

Introduction

Endometriosis is an inflammatory condition characterized by lesions of endometrial-like tissue outside of the uterus and is associated with pelvic pain and infertility (Giudice, 2010). It affects an estimated 176 million women of reproductive age worldwide (Adamson *et al.*, 2010). It is widely assumed that lesions arise through retrograde endometrial tissue loss during menstruation, coelomic metaplasia and lymphatic spread in immunologically and genetically susceptible individuals. While its underlying cause is uncertain, it is likely to be multifactorial including genetic factors with possible epigenetic influences, perhaps promoted through environmental exposures. Endometriosis has elements of a pain syndrome with central neurological sensitization (and some hallmarks of a neurological disorder) (Stratton and Berkley, 2011), and is a proliferative, estrogen-dependent disorder (with growing evidence of progesterone resistance) (Pabona *et al.*, 2012). There is overlap with other conditions characterized by pelvic–abdominal pain and infertility. Some symptomatic women with pelvic pain, who do not have diagnosed endometriosis or who are prior to diagnosis, may benefit from similar treatments.

Women with endometriosis typically have a range of pelvic–abdominal pain symptoms, including dysmenorrhoea, dyspareunia, heavy menstrual bleeding, non-menstrual pelvic pain, pain at ovulation, dyschezia and dysuria, as well as chronic fatigue (Kennedy *et al.*, 2005; Nnoaham *et al.*, 2011). Endometriosis lesions, particularly deep infiltrating lesions, are often innervated. The presence of endometriotic lesions, followed by denervation and re-innervation, may result in accompanying changes in the central nervous system (central sensitization), creating a chronic pain syndrome (Stratton and Berkley, 2011). Endometriosis is also associated with infertility, with a strong association between severity of disease and impact on fertility, probably due to impaired tubo-ovarian function, the presence of ovarian endometrioma, subclinical pelvic inflammation, possibly reduced oocyte quality and reduced endometrial receptivity to implantation (Lessey, 2011). Both endometriosis and adenomyosis (lesions occurring in the uterine intramural muscular layer) reduce the chance of success of assisted reproductive treatment (Barnhart *et al.*, 2002; Maubon *et al.*, 2010).

Symptoms of endometriosis contribute substantially to the burden of disease and add substantial cost to society through reduced economic and personal productivity (Simoens *et al.*, 2007; Nnoaham *et al.*, 2011; Simoens *et al.*, 2012).

While symptoms and examination findings may suggest endometriosis (Nnoaham *et al.*, 2011, 2012), the gold standard for making the diagnosis remains the laparoscopic visualization of lesions preferably with histologic confirmation (Kennedy *et al.*, 2005). In the absence of histological sampling, the false-positive rate with laparoscopic visualization alone may approach 50% especially in women with minimal or mild endometriosis (Wykes *et al.*, 2004). Laparoscopy also enables endometriosis to be staged by the revised American Society for Reproductive Medicine (r-ASRM, 1997) scoring system, the ‘scoring’ system most commonly in current use, objectively defining the disease as minimal (stage I), mild (stage II), moderate (stage III) or severe (stage IV) based on its laparoscopic appearance. It is recognized that the stage/extent of disease may not correlate with symptoms experienced, reproductive outcome or recurrence risk (Adamson, 2011). Much research has recently focused on serum

biomarkers, including cancer antigen-125 (CA125), leptin, monocyte chemoattractant protein-1 (MCP-1), regulated on activation normal T cell expressed and secreted (RANTES) and macrophage migration inhibitory factor (MIF), although these have not been useful diagnostic predictors owing to poor sensitivity or specificity, small sample size or inadequate validation of their accuracy (May *et al.*, 2010). Recent interest has focused on endometrial immunohistochemistry for nerve fibre density (Al-Jefout *et al.*, 2009; Bokor *et al.*, 2009) and on urinary markers (cytokeratin 19, urinary peptide 1.8 kDa) (May *et al.*, 2010). These less invasive diagnostic tests require future formal and robust evaluation of their accuracy.

Several guidelines have been developed by a number of national and international bodies: the European Society for Human Reproduction and Embryology (<http://guidelines.endometriosis.org/>), the American Society of Reproductive Medicine: ([http://www.asrm.org/uploadedFiles/ASRM_Content/News_and_Publications/Practice_Guidelines/Educational_Bulletins/endometriosis_and_infertility\(1\).pdf](http://www.asrm.org/uploadedFiles/ASRM_Content/News_and_Publications/Practice_Guidelines/Educational_Bulletins/endometriosis_and_infertility(1).pdf) and [http://www.asrm.org/uploadedFiles/ASRM_Content/News_and_Publications/Practice_Guidelines/Educational_Bulletins/Treatment_of_pelvic_pain\(1\).pdf](http://www.asrm.org/uploadedFiles/ASRM_Content/News_and_Publications/Practice_Guidelines/Educational_Bulletins/Treatment_of_pelvic_pain(1).pdf)), the Royal College of Obstetricians and Gynaecologists (<http://www.rcog.org.uk/files/rcog-corp/GTG2410022011.pdf>), the Society of Obstetrics and Gynecology of Canada (<http://www.sogc.org/guidelines/documents/gui244CPG1007E.pdf>) and the Cochrane Database of Systematic Reviews (<http://thecochranelibrary.com>), yet areas of controversy and uncertainty remain, not least due to a paucity of firm evidence.

The World Endometriosis Society (WES) has therefore developed a process to bring together representatives of national and international, medical and non-medical societies with an interest in endometriosis, aiming to derive a consensus on the management of endometriosis from a global perspective, in which the views of women with endometriosis were represented.

Methods

We developed a consensus process supported by a specific methodology (Supplementary data, Information 1). This differed from a formal guideline methodology, which typically involves a more lengthy and prescriptive process.

There were 51 national and international societies invited to participate in the WES Consensus on the Management of Endometriosis and to nominate a representative for their organization in the consensus process and at the meeting in Montpellier on 8 September 2011. From these nominees, along with members of the WES Board, a group of participants in the WES Montpellier Consortium was established and this ultimately comprised 56 representatives from 34 organizations (18 medical organizations, 16 non-medical endometriosis organizations). Pharmaceutical companies with an interest in developing products for treating endometriosis were invited to send a representative to the Montpellier meeting as an observer and two companies participated. All participants and their roles are summarized in Table 1.

The participants were involved in an on-going email discussion group for 4 months and a teleconference in advance of the meeting, with the following goals:

- (i) to have all participants conversant with the evidence;
- (ii) to define topics for presentation;
- (iii) to determine reviewers to present these topics;

Table 1 The World Endometriosis Society Montpellier Consortium.

Name	Pre-meeting email consultations	3 August 2011 phone meeting	Presenter	Attended 8 September 2011 meeting	Voted	Manuscript revision
Mauricio Abrao	x	x	x	x	x	x
David Adamson	x	—	x	x	x	x
Catherine Allaire	—	—	—	x	x	x
Vibeke Amelung	—	—	—	—	x	x
Elisabet Andersson	x	—	—	x	x	x
Mary-Lou Ballweg	—	—	—	x	Resigned	
Christian Becker	—	—	—	x	x	x
Kolbrún Birna Árdal	x	—	—	x	—	x
Deborah Bush	x	x	x	x	x	x
Bianca de Bie	—	—	—	x	x	x
Kristof Chwalisz	—	—	—	x	—	x
Hilary Critchley	—	—	—	—	x	x
Thomas D'Hooghe	x	—	x	x	x	x
Gerard Dunselman	x	—	x	x	x	x
Johannes Evers	x	—	—	x	x	x
Cindy Farquhar	x	x	x	x	x	x
Thomas Faustmann	x	—	—	x	x	x
Axel Forman	—	—	—	—	—	x
Jessica Fourquet	—	—	—	x	x	x
Ian Fraser	—	—	—	x	x	x
Linda Giudice	x	x	x	x	—	x
Stephan Gordts	x	—	x	x	x	x
Heather Guidone	x	—	—	—	x	x
Sun-Wei Guo	—	—	—	—	x	x
David Healy	x	—	x	x	Deceased	
Bernard Hedon	—	—	—	x	x	x
Johanna Hulkkonen	x	—	—	x	x	x
Louise Hull	—	—	—	x	x	x
Lone Hummelshoj	x	x	x	x	x	x
Neil Johnson	x	x	x	x	x	x
Miriam Just	x	—	—	x	x	—
Ludwig Kiesel	—	—	—	—	x	x
Alan Lam	—	—	—	—	x	x
Clodagh Lynam	x	—	—	x	x	x
Liselotte Mettler	—	—	—	x	x	x
Charles Miller	x	—	x	x	—	—
Helen North	—	—	—	x	x	x
Rishma Pai	—	—	—	—	x	x
Carlos Petta	x	x	x	x	x	x
Lucy Prentice	—	—	—	x	—	x
Fernando Reis	—	—	—	x	x	x
Shelley Reilly	—	—	—	x	—	x
Edgardo Rolla	x	x	x	x	x	x

Continued

Table I Continued

Name	Pre-meeting email consultations	3 August 2011 phone meeting	Presenter	Attended 8 September 2011 meeting	Voted	Manuscript revision
Luk Rombauts	x	x	x	x	x	x
Karl-Werner Schweppe	x	—	x	x	x	x
Tamer Seckin	x	—	—	x	x	—
Kathy Sharpe-Timms	—	—	—	—	x	x
Dian Shepperson Mills	x	—	—	x	x	x
Sony Singh	x	x	x	x	—	x
David Soriano	—	—	—	x	x	—
Martyn Stafford-Bell	x	—	—	x	x	x
Pamela Stratton	x	x	x	x	x	x
Robert Taylor	x	x	x	x	x	x
Jim Tsaltas	x	—	x	x	x	x
Jacqueline Veit	—	—	—	x	—	x
Paolo Vercellini	x	—	x	x	x	x

Representing: AAGL, Abbott Laboratories, Associazione Italiana Endometriosi, AGES, ALMER, AFOG, American Society for Reproductive Medicine, Bayer Pharma, Cochrane Collaboration, Endometrioseforeningen (Norway), Endometriose Stichting (NL), Endometriose Foreningen (Denmark), Endometrioseforeningen Sweden, Endometriosis Association (USA), Endometriosis Association of Ireland, Endometriosis Foundation of America, Endometriosis New Zealand, Endometriosis Research Center (USA), Endometriosis SHE Trust UK, Endometriosis UK, Endometriosisyhdistys (Finland), European Society of Gynaecologic Endoscopy, European Society of Human Reproduction and Embryology, European Endometriosis Liga, FIGO, Fundación Puertorriquena de Pacientes con Endometriosis, International Federation of Fertility Societies, International Society of Gynecologic Endoscopy, Israeli Endometriosis Society, RANZCOG, Samtök Kvenna med Endómétríósu (Iceland), Sociedade Brasileira de Endometriose, Society of Gynecologic Investigation, Society of Obstetrics and Gynaecology Canada, World Endometriosis Research Foundation, WES.

(iv) to refine the clinical questions addressed, including the patient populations, interventions and outcomes to be considered.

The consensus meeting took place on 8 September 2011 in Montpellier, in association with the 11th World Congress on Endometriosis. Topics were presented by each reviewer, who had been asked to prepare draft consensus statements, based on their extensive literature reviews (see [Supplementary data, Information 2](#)). After full discussion, the proposed consensus statements were modified if necessary by agreement.

The relevant evidence was appraised according to the GRADE system ([Guyatt et al., 2008](#)) (see [Supplementary data, Information 3](#)), leading to a consensus statement, graded as either strong or weak, the abiding principle being that where, across the range of issues considered important by women with endometriosis, fully informed women were likely to make different choices, a weak statement was made ([Guyatt et al., 2008](#)). Where evidence from studies was lacking, but where the group felt that we had sufficient expertise and anecdotal experience to make an important statement, the statement was ascribed a 'good practice point' (GPP) and, through discussion, determined as strong or weak. For GPPs the definition of a strong statement was one where the disease burden was high and the potential impact of an intervention was considerable with minimal downside. The level of consensus around each statement was also ascribed a consensus grade, using the consensus grading system developed by the Australasian CREI Consensus Expert Panel on Trial evidence Group of the Royal Australian and New Zealand College of Obstetricians and Gynaecologists (RANZCOG) ([Kroon et al., 2011](#)). For the consensus grades ascribed to each of our statements, α (unanimous or near-unanimous) was where more than 80% agreed without caveat and fewer than 5% disagreed, β (unanimous with caveat) was where fewer

than 5% disagreed but fewer than 80% agreed without caveat (the major caveats have been highlighted in the text), γ (majority) was where 50–80% agreed and δ (no consensus) was where fewer than 50% agreed with or without caveat.

Evidence tables ([Supplementary data, Information 4](#)) were established for all the evidence considered at the consensus meeting. Where the evidence base was considered to be well established, for example with medical treatments for endometriosis (first and second line), the evidence was amalgamated into a single table. For treatments where the evidence base was less well established, particularly for emerging treatments or complementary therapies, each table represented evidence for individual treatments.

A consensus statement was drafted by the meeting conveners (N.J. and L.H.) with further reference to the Power Point presentations and an audiotape of the proceedings of the meeting. A post-meeting online survey was conducted to systematically define the consensus around each of the statements made by a formal voting procedure. Of the WES Montpellier Consortium, 57% ($n = 32$) contributed to the pre-meeting debates, 84% ($n = 47$) attended the meeting in Montpellier and 80% ($n = 45$) completed the post-meeting online survey. One participant was deceased and one participant resigned from the Consortium after the meeting. Those contributing to this consensus, who did not attend the meeting in Montpellier, acted as first-level external reviewers ($n = 9$). Following three rounds of modification by circulation to and feedback from the WES Montpellier Consortium, the consensus statement was finalized. A more detailed version of the methodology may be found in [Supplementary data, Information 1](#). Consortium members' contributions at each step of the process are outlined in [Table 1](#).

Results

The WES consensus statements

The evidence tables ([Supplementary data, Information 4](#)) provide the evidence that was considered to reach the consensus statements. The consensus statements, categorized as either strong or weak, are summarized in [Table II](#), along with the level of consensus that applied to each statement.

General principles

It was suggested that a philosophical shift to consideration of ‘endometriosis and pelvic pain’ as a spectrum or continuum of disease will avoid excluding women who lack laparoscopic confirmation of a diagnosis of endometriosis.

Endometriosis in low-resource settings

From a global perspective, there was strong consensus that diagnosis and management of endometriosis should be incorporated into the primary health care of women worldwide. In low-resource settings, diagnosis may commence with two simple questions about pelvic-abdominal pain and infertility (accepting that a negative response does not exclude endometriosis). Management, including prevention, should be integrated with other women’s healthcare strategies in low-resource settings, and may include education, progestin-based contraceptives, family planning and lactation.

Centres/networks of expertise

Women with endometriosis often require individualized care over a long-term period, where priorities may change depending upon the type and severity of symptoms, impact of these symptoms, current or future fertility goals and lifestyle factors. However, not all women with endometriosis require a large number of experts and some women are treated effectively for the rest of their lives by a single laparoscopic surgical procedure. Individualized care benefits from a multi-disciplinary network of experts sufficiently skilled in providing advice on and treatment of endometriosis and its associated symptoms, based on the best available evidence, their extensive experience and their transparent record of success rates. Previously the term ‘centre of excellence’ has been used ([D’Hooghe and Hummelshoj, 2006](#)) but we now agree that ‘centre (or network) of expertise’ is more appropriate. It was accepted that a centre/network of expertise would take differing forms in different settings, although consensus over precisely what form this would take (involving either a team, a network or a physical unit or centre where expertise is concentrated and coordinated) was not reached. However, it was agreed that such centres/networks should ideally comprise a multi-disciplinary team approach with specialists who have undergone specific training in endometriosis, advanced surgeons with a high caseload of managing deep endometriosis (also known as deep infiltrating endometriosis, DIE), ready access to an endometriosis organization with substantial input on behalf of women and a track record of commitment to collaborative management and research. As laparoscopic surgery will likely continue to be pivotal in the management of women with endometriosis, accreditation should be focused on the training and expertise of laparoscopic surgeons. The centre/network should have a transparent record of outcome-based success rates. There was no

consensus on the accreditation or longevity of such an accreditation. Whilst it is impractical that all women with endometriosis are currently managed in a centre/network of expertise, those with higher stage of disease and/or more intractable clinical problems should receive care from such a centre or network.

Endometriosis support groups and endometriosis organizations

National endometriosis support groups and endometriosis organizations exist around the world. Feedback from women and endorsement from health professionals and other stakeholders substantiate the value of effective support groups and endometriosis organizations to individuals ([Kennedy et al., 2005](#); [Bush, 2009](#)), although not all women need these services. Endometriosis support groups provide a valuable forum for women with endometriosis, having the potential to assist women to improve their quality of life by teaching coping mechanisms and sharing experiences. Engagement of experienced and skilled medical practitioners, accredited educators and other stakeholders brings strength to an endometriosis organization.

Life journey of women with endometriosis

The stage of a woman’s life is an important determinant of her requirement for treatment options, particularly according to her current symptoms, including present or future fertility wishes. Most of the consensus statements that follow relate to women within the reproductive age group; however it is acknowledged, as follows, that endometriosis may persist after natural or surgical menopause and must be managed accordingly.

Adolescents with endometriosis

Endometriosis should be considered as a possible diagnosis in adolescents with suggestive symptoms—most women diagnosed with endometriosis date the onset of their symptoms to their teens ([Nnoaham et al., 2011](#)). Most adolescents have stage I or II disease ([Laufer et al., 1997](#)), although endometriosis of any stage may present in adolescence ([Roman, 2010](#)). Currently, there is insufficient evidence to make strong recommendations for management amongst adolescents who may have endometriosis ([Dovey and Sanfilippo, 2010](#); [Yeung et al., 2011](#)). Treatment (both medical and surgical) for this age group may improve the quality of life, reduce symptoms, prevent more severe disease developing later and reduce the likelihood of compromised future fertility, but further research to clarify these issues is essential. An appropriate balance of discussion of endometriosis as a possible diagnosis, then appropriate treatment (either empirical medical or surgical), without an over-interventional approach, must be sought. There is a pressing need for research into and guidelines for the management of symptomatic endometriosis and possible endometriosis amongst adolescents.

Obstetric outcomes for women with endometriosis

Evidence is emerging that women with endometriosis have a higher risk of obstetric complications, including preterm delivery, antepartum haemorrhage, possibly pre-eclampsia and Caesarean section ([Fernando et al., 2009](#); [Stephansson et al., 2009](#); [Brosens et al., 2012](#)), in addition to rare life-threatening situations where intra-abdominal bleeding from endometriotic lesions can lead to the need for urgent

Table II World Endometriosis Society Montpellier Endometriosis Consensus Statements.

	Consensus grading
Endometriosis in low-resource settings	α
(1) Endometriosis diagnosis and management should be incorporated into the primary health care of women worldwide (strong GPP).	
(2) In low-resource settings, diagnosis may commence with two simple questions about pelvic–abdominal pain and infertility (strong GPP).	β
(3) Management, including prevention, should be integrated with other women’s healthcare strategies in low-resource settings, and may include education, progestin-based contraceptives, family planning and lactation (strong GPP).	α
Networks of expertise	
(4) Women with endometriosis require individualised care over a long-term period, where priorities may change owing to the type and severity of symptoms, impact of these symptoms, current or future fertility wish and lifestyle factors (strong GPP).	α
(5) Individualised care benefits from a multi-disciplinary network of experts sufficiently skilled in providing advice on and treatment of endometriosis and its associated symptoms, based on the best available knowledge, their extensive experience and their transparent record of success rates (strong GPP).	β
Endometriosis organizations and support groups	
(6) Endometriosis support groups provide a valuable forum for women with endometriosis having the potential to assist women to improve their quality of life by teaching coping mechanisms and sharing experiences (strong GPP).	γ
(7) Engagement of experienced and skilled medical practitioners, accredited educators and other stakeholders brings strength to an endometriosis organization (strong GPP).	α
(8) A philosophical shift to consideration of ‘endometriosis and pelvic pain’ as a spectrum or continuum of disease will avoid excluding women who lack laparoscopic confirmation of a diagnosis of endometriosis (weak GPP).	γ
Endometriosis and adolescence	
(9) Endometriosis should be considered as a possible diagnosis in adolescents with suggestive symptoms (strong).	α
(10) Currently, there is insufficient evidence to make strong recommendations for management amongst adolescents who may have endometriosis (weak).	γ
Endometriosis and obstetric outcomes	
(11) Endometriosis should be considered an obstetric risk factor and pregnancies managed accordingly (strong).	γ
Endometriosis and menopause	
(12) Although endometriosis may occasionally recur, there is no strong evidence to deprive women of HRT if they suffer severe menopausal symptoms but have a history of endometriosis, although combined estrogen-progestin hormone therapy is advisable (weak).	γ
Endometriosis and cancer	
(13) The relative risk and absolute risk of ovarian cancer amongst women with endometriosis is so low as not to justify routine ovarian cancer screening (strong).	γ
Lifestyle/dietary interventions	
(14) Dietary intervention following endometriosis surgery in the form of vitamins, minerals, salts, lactic ferments and fish oil appears to be a suitable alternative to hormonal treatment, that is associated with similar pelvic pain reduction and quality of life improvement (weak).	δ
Empirical medical treatment	
(15) Well-tolerated, low-cost, easily accessible options such as non-steroidal anti-inflammatory drugs (NSAIDs), other analgesics, combined OCP and progestins should be considered for use as first-line empirical medical treatment (strong).	γ
(16) In some circumstances, second-line medical treatment with gonadotrophin-releasing hormone agonists (GnRH-a) with add-back HRT, or the LNG-IUS may be considered for use as empirical medical treatment for women who are not optimally treated with first-line empirical therapy prior to surgical diagnosis and treatment, whilst awaiting laparoscopic surgery (weak).	γ
Surgery for women with symptomatic endometriosis	
(17) Laparoscopic surgical removal of endometriosis is an effective first-line approach for treating pain related to endometriosis (strong).	α
(18) Although current RCTs have failed to demonstrate benefit of excision over ablation, it is recommended to excise lesions where possible, especially deep endometriotic lesions (weak).	α
(19) Laparoscopic surgery for endometriosis should always be undertaken in preference to laparotomy, where possible (strong GPP).	γ
(20) The addition of LUNA to laparoscopic removal of endometriosis does not improve pain relief (strong).	β
(21) Although PSN might benefit a small number of women, the benefits are likely to be outweighed by the potential for harmful effects (strong).	γ
(22) Laparoscopic excision (cystectomy) for ovarian endometriomas is preferred where possible to minimise symptom recurrence and endometrioma recurrence (strong).	γ
(23) The best surgical approach to deep endometriosis is unclear (weak).	γ

Continued

Table II *Continued*

	Consensus grading
(24) Highly specialised surgical expertise is required by surgeons, who undertake surgery for deep endometriosis, and it should be undertaken only within centres of expertise (strong GPP).	α
Medical therapy for women with symptomatic endometriosis	
(25) Well-tolerated, low-cost, easily accessible options such as non-steroidal anti-inflammatory drugs (NSAIDs), other analgesics, combined OCP and progestins should be considered for first-line medical treatment of laparoscopically diagnosed endometriosis (strong).	γ
(26) The combined OCP is an effective medical treatment to minimise the endometrioma recurrence rate after surgical removal of the cyst (strong).	α
(27) Second-line medical treatments could include gonadotrophin-releasing hormone agonists (GnRH-a, which should be used with add-back HRT, routinely), the LNG-IUS and depot progestins (weak).	γ
(28) Danazol and gestrinone should not be used other than for women, established on these treatments in the absence of side effects, for whom other treatments have proven ineffective (strong).	α
Emerging medical therapies for women with symptomatic endometriosis	
(29) Aromatase inhibitors might be reasonable as a second-line medical treatment, but more research is required (weak).	γ
(30) SPRMs might be a reasonable second-line medical treatment, but more research is required (weak).	γ
(31) Gonadotrophin-releasing hormone (GnRH) antagonists might be reasonable as second-line medical treatment, but more research is required (weak).	γ
(32) There is no evidence of a benefit of pentoxifylline on the reduction of pain (strong).	α
(33) There is no evidence of a benefit of anti-TNF α (anti tumour necrosis factor alpha) on the reduction of pain (weak).	γ
(34) There is no benefit from raloxifene on prevention of recurrence of pain (strong).	α
(35) There is insufficient evidence of a benefit of rosiglitazone on the reduction of pain (weak).	γ
(36) There is insufficient evidence of benefit of valproic acid on the reduction of pain (weak).	γ
(37) Anti-angiogenesis agents are at research level only (strong).	α
Complementary therapies for women with symptomatic endometriosis	
(38) There is some evidence of effectiveness of acupuncture, but it requires repeated treatments and effects are unlikely to be long lasting (weak).	γ
(39) There is evidence of effectiveness of TENS for short-term pain management for women with dysmenorrhoea (weak).	γ
(40) There is insufficient evidence of effectiveness of traditional Chinese medicine (TCM) and applicability is uncertain outside of TCM settings (weak).	α
(41) Vitamin B1 and B6 can be used to relieve pain for women with dysmenorrhoea but there is limited evidence of effectiveness and there are safety concerns with vitamin B6 at higher doses (weak).	γ
(42) There is some evidence of effectiveness of magnesium in reduction of pain for women with dysmenorrhoea (weak).	γ
(43) There is no evidence of effectiveness for topical heat (weak).	γ
(44) There is no evidence to support spinal manipulation (weak).	γ
(45) There is insufficient evidence to support behavioural interventions (weak).	γ
Surgery for infertility in women with endometriosis	
(46) Laparoscopic surgical removal of endometriosis improves fertility in stage I and II endometriosis (strong).	γ
(47) Although RCTs have failed to demonstrate benefit of excision over ablation, it is recommended to excise lesions where possible, especially where pain is present (weak).	γ
(48) Laparoscopic excision (cystectomy) where possible for endometriomas is preferred to laparoscopic ablation (drainage and coagulation) to enhance fertility (strong).	α
(49) The best surgical approach to deep endometriosis in women with infertility is unclear (weak).	γ
(50) Medical adjunct therapy in conjunction with laparoscopic surgery has not been shown to have fertility benefit (strong).	α
Assisted conception for infertility in women with endometriosis	
(51) There is no evidence to support the use of controlled OS alone and insufficient evidence to recommend one agent over another (weak).	γ
(52) Intrauterine insemination (IUI) with controlled OS (COS) is effective in improving fertility in minimal and mild endometriosis, but the role of unstimulated IUI is uncertain (strong).	γ
(53) Double insemination should be considered for intrauterine insemination (IUI) (weak).	δ

Continued

Table II *Continued*

	Consensus grading
(54) Although IVF may be less effective for endometriosis than for other causes of infertility, it should be considered for use to improve the success rate above expectant management (strong).	γ
Adjuncts to assisted conception for infertility in women with endometriosis	
(55) There is insufficient evidence of benefit of gonadotrophin-releasing hormone (GnRH-a) treatment before intrauterine insemination (IUI) (weak).	α
(56) There is insufficient evidence of benefit of laparoscopic surgery prior to IUI/COS (weak).	γ
(57) GnRH analogue administered for 3–6 months prior to IVF/ICSI in women with endometriosis increases the clinical pregnancy rate (strong).	γ
(58) There is insufficient evidence to support the use of the combined OCP prior to IVF/ICSI (weak).	γ
(59) There are no data to compare the approach of pretreatment with the combined OCP versus gonadotrophin-releasing hormone agonists (GnRH-a) (weak).	γ
(60) There is no evidence that surgical removal of endometriosis or surgical treatment of endometriomas (by aspiration or cystectomy) improves success rates through IVF (weak).	γ
(61) Ovarian response might be reduced in some women who have undergone surgery for endometriomas (weak).	α
(62) Since endometriomas may damage the ovary, and since complications can arise in women with endometriomas undergoing ART, laparoscopic ovarian cystectomy may sometimes be recommended for women with endometriomas larger than 3 cm diameter (weak).	α
Medical therapy for infertility in women with endometriosis	
(63) There is no evidence of fertility benefit from medical treatment—ovulation suppression may delay pregnancy and this is not recommended (strong).	α
Emerging therapies for infertility in women with endometriosis	
(64) Lipiodol hysterosalpingogram improves live birth rates in women with endometriosis, but otherwise unexplained infertility, who are attempting natural conception (weak).	γ
(65) There is no evidence of fertility benefit from pentoxifylline for women with mild-to-moderate endometriosis (strong).	α
(66) There is no evidence of fertility benefit of TCM over gestrinone or Danazol (weak).	γ
(67) There is insufficient evidence of increased pregnancy rates from the use of vitamins (weak).	α
(68) There is insufficient reliable evidence of improved fertility with mifepristone (weak).	α
(69) There is no evidence of impact of rosiglitazone on fertility (weak).	α

The above represent the consensus statements from the WES Montpellier Consensus.

GPP, good practice point; α , unanimous or near-unanimous (more than 80% agreed without caveat and fewer than 5% disagreed); β , unanimous with caveat (fewer than 5% disagreed but fewer than 80% agreed without caveat); γ , majority (50–80% agreed); δ , no consensus (fewer than 50% agreed with or without caveat).

surgery (Mutahir and Nyango, 2010). It was agreed that a history of endometriosis should be considered an obstetric risk factor and pregnancies managed accordingly.

Menopausal women with endometriosis

It has been reported that after a diagnosis of endometriosis, 96.9% of women become free from pain after menopause (Fagervold et al., 2009). However, post-menopausal endometriosis has seldom been investigated, though symptoms usually disappear after a natural or a surgical menopause. The risk of recurrence with hormone therapy is probably increased in women with residual disease after surgery and the consequent management is best monitored by responding to specific symptoms (Moen et al., 2010). Although endometriosis may recur, there is no strong evidence to deprive women of hormone replacement treatment (HRT) if they suffer severe menopausal symptoms but have a history of endometriosis, although combined estrogen-progestin hormone therapy is advisable (Al Kadri et al., 2009; Moen et al., 2010).

Ovarian cancer

There is a recognized association between endometriosis and clear cell, low-grade serous and endometrioid ovarian cancer (Pearce et al., 2012), but the overall risk of ovarian cancer amongst women with endometriosis remains low, with a relative risk ranging from 1.3 to 1.9 (Sayasneh et al., 2011) which means that at worst the lifetime risk of ovarian cancer is increased from ~1 in 100 to 2 in 100. Yet so far there is no unequivocal evidence of causality in this association. Thus, there is no evidence in favour of routine ovarian cancer screening for women with endometriosis, but the question remains as to whether there is a higher risk group amongst women with endometriosis in whom screening may be justified (such as those with recurrent ovarian cysts or suspected but unremoved endometrioma in the menopause). It is recommended that future studies must endeavour to clearly establish or exclude causality rather than mere association due to shared risk factors. Establishing a genetic basis of subgroups of women with endometriosis may lead to the identification of any pre-disposition of certain cancers, and thus a possible identification of high-risk subgroups. Only then can specific clinical guidelines be recommended.

Lifestyle and dietary interventions

Whilst the overwhelming response from women managing their endometriosis is that these interventions do help to improve the quality of life, few well-designed studies have examined lifestyle factors. Examples of lifestyle interventions described as helpful but not so far exposed to randomized controlled trial (RCT) scrutiny include simply 'talking to someone', cognitive behavioural therapy and different types of exercise including yoga. No well-designed studies have examined exercise, but a small observational retrospective study suggests that exercise might be effective in reducing dysmenorrhoea (Koppan *et al.*, 2010). There is no evidence to support weight reduction having a beneficial impact on symptomatic endometriosis. No consensus could be established regarding dietary interventions, although evidence from two RCTs showed that dietary intervention following endometriosis surgery in the form of vitamins, minerals, salts, lactic ferments and fish oil appears to be an effective alternative to hormonal treatment, that is associated with similar pelvic pain reduction and quality of life improvement (Sesti *et al.*, 2007, 2009). Observations that certain diets (especially a gluten-free diet) improve symptoms for some women with endometriosis remain unconfirmed in RCTs. For dysmenorrhoea in the absence of proven endometriosis, one small trial showed fish oil (omega-3 fatty acids) to be more effective than placebo for pain relief (Proctor and Murphy, 2001).

Empirical medical treatment for symptoms of endometriosis

Many clinicians support empirical medical treatment of endometriosis either prior to or without laparoscopic confirmation of endometriosis. Time to surgery may delay appropriate treatment, there is a false-negative rate in laparoscopic diagnosis, and surgery is invasive and expensive compared with empirical therapies, and carries a risk of morbidity. Nonetheless, a full evaluation that includes consideration of other causes of the symptoms and assessment of the disease impact for the woman is required prior to empirical treatment. Management of pelvic pain should not be delayed in order to obtain surgical confirmation of endometriosis, even though most of the RCT evidence is from women with surgically confirmed endometriosis. Although the definition of medical treatments as first line versus second line is arbitrary, we adopted as first line those treatments that most clinicians would consider using empirically and second line those treatments that most would reserve for treatment following laparoscopic diagnosis. Well-tolerated, low-cost, easily accessible options such as non-steroidal anti-inflammatory drugs (NSAIDs) (Allen *et al.*, 2009), other analgesics (paracetamol and opioids, although most clinicians would reserve opioid analgesics for second-line treatment), the combined oral contraceptive pill (OCP) (Davis *et al.*, 2007; Harada *et al.*, 2008; Guzik *et al.*, 2011; Vercellini *et al.*, 2011) and traditional progestins such as medroxyprogesterone acetate (Crosignani *et al.*, 2006; Schlaff *et al.*, 2006) and norethisterone (Vercellini *et al.*, 2011; Brown *et al.*, 2012) or newer progestins such as dienogest (Cosson *et al.*, 2002; Harada *et al.*, 2009; Momoeda *et al.*, 2009; Köhler *et al.*, 2010; Strowitzki *et al.*, 2010a, b, 2012; Petraglia *et al.*, 2012), should be considered for use as first-line empirical medical treatment. Some clinicians would, in certain circumstances, consider second-line medical treatment with gonadotrophin-releasing hormone agonists (GnRH-a) with add-back HRT (Brown *et al.*, 2010), the levonorgestrel-releasing intrauterine system (LNG-IUS) (Abou-Setta

et al., 2006) or opioid analgesics as empirical treatment for women who are not optimally treated with first-line empirical therapy prior to surgical diagnosis and treatment, whilst awaiting laparoscopic surgery (and some women successfully treated with second-line empirical medical treatment might not proceed to surgery). It is unclear whether medical treatment prior to laparoscopy might mask the diagnosis by reducing the appearance of endometriotic implants and hence may make endometriosis more difficult to treat surgically. It is important to highlight that NSAIDs have important side effects, including peptic ulceration and an adverse impact on ovulation, and that analgesics, particularly opiates, if used inappropriately and without medical monitoring, carry a risk of abuse and/or addiction. All women receiving medical treatment should be carefully monitored with regular follow-up consultations.

Surgical management of endometriosis symptoms

The issue of appropriate laparoscopic surgical training is considered vital and there are strong arguments for standardization of what constitutes the relevant experience and expertise for those undertaking complex laparoscopic surgery for endometriosis. Crucial aspects in planning laparoscopic surgery are that surgery should be carried out in the most appropriate setting which can ensure adequate preoperative counselling, appropriate surgical expertise (to ensure the most appropriate procedure is undertaken by the most experienced surgeon at the most appropriate time), adequate technical resources and post-operative support care. Whenever possible, laparoscopic surgery should always be undertaken in preference to laparotomy. It is also important, particularly in cases of more severe endometriosis, that surgeons consider the option of limiting surgical excision at an initial operation in order to refer to a surgeon better equipped to deal with endometriosis, as the first definitive surgical intervention has been shown to deliver the greatest benefit (Abbott *et al.*, 2004).

Laparoscopic surgical removal of endometriosis (through either excision or ablation of endometriosis or both) is an effective first-line approach for treating pain related to endometriosis (Jacobson *et al.*, 2009). Although RCTs have failed to demonstrate the benefit of excision over ablation (Wright *et al.*, 2005; Healey *et al.*, 2010), there is unanimous consensus over the recommendation to excise lesions where possible, especially deep endometriotic lesions, which is felt by most surgeons to give a more thorough removal of disease (Koninckx *et al.*, 2012). It is also acknowledged that, even after expert removal of endometriosis, there may be a recurrence rate of symptoms and endometriotic lesions that varies from 10 to 55% within 12 months (Vercellini *et al.*, 2009), with recurrence affecting ~10% of the remaining women each additional year (Guo, 2009). The risk of requirement for repeat surgery is higher in women younger than 30 years at the time of surgery (Shakiba *et al.*, 2008). First operations tend to produce a better response than subsequent surgical procedures, with pain improvements at 6 months in the region of 83% for first excisional procedures versus 53% for second procedures (Abbott *et al.*, 2004). Excessive numbers of repeat laparoscopic procedures should therefore be avoided. The role of a purely diagnostic laparoscopy has been questioned and, ideally, there should always be the option of continuing to surgical removal of endometriosis, within the limitations of the surgeon's expertise.

There is insufficient evidence to necessitate the planning of surgery for a particular time of the cycle; however, surgery in the follicular

phase avoids the complicating factor of the presence of a haemorrhagic corpus luteum and one study suggested an increased recurrence rate for surgery undertaken in the luteal phase, hypothesized to be due to re-implantation through retrograde loss of endometrial tissue at subsequent menses whilst the sites of surgically removed lesions were still healing (Schweppe and Ring, 2002).

There is no place for adding laparoscopic uterine nerve ablation (LUNA) to laparoscopic removal of endometriosis (Proctor *et al.*, 2005). Although presacral neurectomy (PSN) might provide benefit for a small number of women with central dysmenorrhoea, the benefits are likely to be outweighed by the potential for harmful effects (including presacral haematoma and dysfunction of bladder and/or bowel) and PSN is not usually recommended (Proctor *et al.*, 2005). Laparoscopic PSN, if ever undertaken, should be performed only by expert surgeons.

Laparoscopic excision (cystectomy) for ovarian endometriomas is preferred to laparoscopic ablation (drainage and coagulation) where possible to minimize symptom recurrence and endometrioma recurrence, although care must be taken to minimize damage to surrounding normal ovarian tissue (Hart *et al.*, 2008). Despite most endometriotic cysts being predominantly extra-ovarian in nature, systematic cystectomy performed by highly experienced surgeons has been shown to reduce ovarian volume (Biacchiardi *et al.*, 2011). The value of a multiple-step procedure (interval surgery that utilizes intervening medical suppressive treatment) requires further evaluation, particularly for large ovarian endometriomas (Tsolakidis *et al.*, 2010).

Although the OCP reduces the recurrence rate of endometriomas after ovarian cystectomy (Seracchioli *et al.*, 2010), evidence does not otherwise support the use of short-term pre- or post-operative medical treatment, in association with laparoscopic removal of endometriosis, for improving pain outcomes or recurrence rates (Furness *et al.*, 2009).

Different approaches have been taken to surgery for deep endometriosis. The dilemma is that incomplete resection may reduce symptomatic outcomes (Vercellini *et al.*, 2006), but that radical interventions increase the risk of major complications such as ureteric and rectal injuries (Koninckx *et al.*, 1996). Evidence is still lacking to guide the best surgical approach to deep endometriosis. If the disease includes bowel endometriosis, the surgical options for the bowel include shaving, disc excision or segmental excision and re-anastomosis. Rather than undertake bowel surgery initially, the optimal approach is to first consider medical treatment. Bowel surgery should only proceed on the basis of shared decision-making after thorough consideration of risks versus benefits, ideally following multi-disciplinary consultations that include provision of information for women on potential complications of surgery. Only then should bowel surgery be performed laparoscopically by experts, avoiding laparotomy whenever possible. What is clear is that highly specialized surgical expertise is required in surgery for deep endometriosis and it should be undertaken only in centres of expertise.

Debate continues over the role of hysterectomy and of concurrent oophorectomy, with little reliable evidence to inform practice, but if such surgery is undertaken, it should be performed laparoscopically where possible. Observational studies have suggested improved pain outcomes for women who undergo hysterectomy for r-ASRM Stage IV endometriosis (Ford *et al.*, 2004), but this may be related to associated pathology such as adenomyosis.

Medical management of endometriosis symptoms

We again arbitrarily defined as first line those medical treatments that most clinicians would consider using empirically and second line those treatments that most would reserve for treatment following laparoscopic diagnosis. Medical treatment may be given routinely as an adjunct to surgery either pre- or post-operatively (see above under surgical management), as a defined course of treatment remote from surgery or as a longer term medical treatment strategy designed to prevent recurrence of endometriosis or endometriomas (Vercellini *et al.*, 2013).

Well-tolerated, low-cost, easily accessible options such as NSAIDs (Allen *et al.*, 2009), other analgesics (that include paracetamol, with an aim of effective pain relief) and OCPs can be considered for use as first-line medical treatment of laparoscopically proven endometriosis (Davis *et al.*, 2007; Harada *et al.*, 2008; Guzick *et al.*, 2011; Vercellini *et al.*, 2011); OCPs are particularly effective in minimizing endometrioma recurrence rates after surgical removal of the cyst (Seracchioli *et al.*, 2010). Progestins with a proven effect in RCTs and with a specific indication for the treatment of endometriosis such as medroxyprogesterone acetate (Crosignani *et al.*, 2006; Schlaff *et al.*, 2006), norethisterone (Vercellini *et al.*, 2011; Brown *et al.*, 2012) and dienogest (Cosson *et al.*, 2002; Harada *et al.*, 2009; Momoeda *et al.*, 2009; Köhler *et al.*, 2010; Strowitzki *et al.*, 2010a, b, 2012; Petraglia *et al.*, 2012) can also be considered as first-line treatments taking into consideration their different side-effect profiles. It is important to discuss potential side effects with the woman before treatment commences, and careful monitoring through regular follow-up appointments is recommended.

Second-line medical treatments could include GnRH-a (Brown *et al.*, 2010), which should be used with add-back HRT routinely (Farmer *et al.*, 2009), LNG-IUS, despite more research into effectiveness and relative effectiveness being required (Abou-Setta *et al.*, 2006), depot progestins, although the side-effect profile and thus treatment burden is high (Bayoglu *et al.*, 2011), and opioid analgesics. Other possible second-line medical treatments include non-oral combined hormonal contraceptives, such as transdermal patches and vaginal rings (Vercellini *et al.*, 2010). Danazol and gestrinone should not be used owing to the high-treatment burden of androgenic side effects (Selak *et al.*, 2007), other than for women, established on these treatments in the absence of side effects, for whom other treatments have proven ineffective. Again, acceptable side effects need to be discussed carefully with the woman.

Hypothetically, medical maintenance therapy might be an effective treatment option that could, in some cases, control the denervation and re-innervation changes that are believed to precede central sensitization and the development of a chronic pain syndrome. Whilst the use of medical treatments such as OCP may be long term, specific studies are needed to investigate whether medical intervention may prevent the development of a chronic pain syndrome. However, most medical agents are only effective for the duration of their use and symptoms often recur on treatment cessation.

Emerging medical treatments for management of endometriosis symptoms

For the emerging medical treatments, data are insufficient to recommend these treatments for routine clinical use. Aromatase inhibitors

(anastrozole, fadrozole, formestane, exemestane, letrozole) (Ferrero *et al.*, 2011), selective progesterone receptor modulator (mifepristone, ulipristal) (Guo *et al.*, 2011) and orally active GnRH antagonists (elagolix) (Struthers *et al.*, 2009) have shown some promise and effectiveness in RCTs, but more clinical experience is required with these agents and more clinical trial research data are essential, especially with regard to their long-term efficacy and side effects. For the immunomodulator, pentoxifylline (Lu *et al.*, 2012), and the anti-TNF- α agent, infliximab (Koninckx *et al.*, 2008), RCTs have not shown benefit to date. The selective estrogen receptor modulator (SERM), raloxifene, has been shown not to provide benefit (Stratton *et al.*, 2008). Possible future treatments yet to be exposed to RCT scrutiny, but where observational studies and case series have suggested promise, include the selective progesterone receptor modulators (SPRMs) asoprisnil and megestrol (Spitz, 2009), the thiazolidinedione, rosiglitazone (Moravek *et al.*, 2009) and valproic acid (Liu and Guo, 2008). As angiogenesis is a crucial activity for the normal processes of the reproductive tract and other organ systems, it is dubious whether agents used for their anti-angiogenic properties (including cabergoline, endostatin, sirolimus, thalidomide and vascular endothelial growth factor inhibitors) will be useful clinically and these have been used only in animal research to date (Laschke and Menger, 2012).

Complementary therapies for endometriosis symptoms

Complementary therapies may help women to cope better with their endometriosis and its treatment and are supported by some evidence from RCTs.

Endometriosis specific. Acupuncture appears to be moderately effective and safe but requires repeated treatments (Zhu *et al.*, 2011). High-frequency transcutaneous electrical nerve stimulation (TENS) has some effectiveness for short-term pain management (Proctor *et al.*, 2002). There is limited evidence in favour of Chinese herbal medicine that may be difficult to apply outside of the Traditional Chinese Medicine setting (Zhu *et al.*, 2008; Flower *et al.*, 2012). While a voluminous literature exists in almost exclusively Chinese medical journals, various problems in study design, execution, statistical analysis and reporting among papers published in Chinese journals make it extremely difficult to judge the efficacy of the evaluated herbal medicine (Guo *et al.*, 2010).

Dysmenorrhoea only. There is limited evidence of benefit for vitamins B1 and B6, with safety concerns associated with higher doses of vitamin B6 (Proctor and Murphy, 2001). Moderate quality evidence supports the use of magnesium (Proctor and Murphy, 2001). Topical heat may be effective for low back pain, but there are no studies specifically examining dysmenorrhoea (French *et al.*, 2006). Spinal manipulation (Proctor *et al.*, 2006) and behavioural interventions (Proctor *et al.*, 2007) are not recommended currently, with more research required for these types of interventions. Cannabis has been shown to be moderately effective for relieving chronic pain (Lucas, 2012), but its benefits are far outweighed by potentially serious side effects and there are no studies in women with endometriosis.

Surgery for endometriosis-associated infertility

The principles of laparoscopic surgery for subfertility are similar to those for other endometriosis symptoms. Appropriate surgical training

is again the key to the best outcomes. It is very important to consider ovarian reserve prior to laparoscopic surgery in the woman experiencing infertility (Pellicano *et al.*, 2008) in particular because evidence is growing that surgical treatment of endometriomas contributes to reduced ovarian reserve (Somigliana *et al.*, 2012; Streuli *et al.*, 2012). The co-existence of pain will be an important factor to consider that will impact on the decision whether to proceed with surgery, although surgery and ART should be considered as complementary strategies.

Laparoscopic surgical removal of endometriosis is recognized as being effective in improving fertility in stage I and II endometriosis (Jacobson *et al.*, 2010). Although RCTs have failed to demonstrate the benefit of excision over ablation, it is recommended to excise lesions where possible, especially deep endometriosis where pain is present (Koninckx *et al.*, 2012). No RCTs have to date assessed whether surgery improves fertility in stage III and IV endometriosis and in deep endometriosis. The functional appearance of the fallopian tubes and ovaries at the end of the laparoscopic procedure appears to contribute to the chance of natural conception post-operatively (Adamson and Pasta, 2010).

Laparoscopic excision (cystectomy) whenever possible for endometriomas >4 cm in diameter improves fertility more than ablation (drainage and coagulation) (Hart *et al.*, 2008). However, much care needs to be taken in identification of tissue planes and careful dissection of the endometrioma to avoid removing normal ovarian tissue and thus impacting on ovarian reserve. There is also the possibility that suturing for haemostasis might maintain ovarian reserve more effectively than electrosurgical haemostasis (Pellicano *et al.*, 2008) and, at the very least, minimization of the use of energy modalities in haemostasis is imperative. Young women, for whom fertility is a consideration, might benefit from discussion of the option of oocyte freezing prior to undergoing ovarian endometrioma surgery, especially if bilateral.

The best surgical approach for deep endometriosis in the context of endometriosis-related infertility remains unclear, even though observational studies suggest good fertility results in women who undergo laparoscopic excision (Chapron *et al.*, 1999; Vercellini *et al.*, 2006; Barri *et al.*, 2010) or laparoscopic shaving (Donnez and Squifflet, 2010). Similarly, colorectal excision is suggested to be beneficial in observational studies (Ferrero *et al.*, 2009; Stepniewska *et al.*, 2010). So far these surgical approaches have not been assessed in RCTs and carry a high risk of complications. Laparoscopic surgery for deep endometriosis, including colorectal endometriosis, should be considered a second-line treatment after failed IVF (unless IVF is not feasible or the patient has severe pain symptoms) and its place in the absence of on-going pain symptoms needs further evaluation.

The pregnancy rate after repeat surgery is lower, approximately half that after first surgery (Vercellini *et al.*, 2009), and two cycles of IVF might be more effective, but surgery should be considered for women with endometriosis-related infertility who continue to be symptomatic or have enlarging endometriomas, and women for whom IVF is declined or repeatedly unsuccessful.

Medical adjunct therapy in conjunction with laparoscopic surgery has not been shown to benefit fertility and is not recommended (Furness *et al.*, 2009); post-operative medical adjunct therapy may delay pregnancy at a time when fertility has been improved by surgery.

Assisted conception for endometriosis-associated infertility

In terms of medically assisted reproduction (MAR), IUI combined with ovarian stimulation (OS) is an effective option for women with minimal-to-mild endometriosis, if the fallopian tubes are normal (Tummon *et al.*, 1997; Costello, 2004). IUI/COS is more effective than unstimulated IUI, with gonadotrophin stimulation appearing to be more effective than that with clomiphene, and the role of unstimulated IUI is uncertain for women with endometriosis (Costello, 2004). However, multiple pregnancy is a key hazard of OS and all reasonable steps should be employed to avoid multiple pregnancy. No consensus could be established over double insemination for IUI (Subit *et al.*, 2011). However, IVF is commonly offered first line in preference to IUI when endometriosis is more severe and tubal function is impaired, or in the context of advanced female age and/or reduced sperm quality.

It is unclear whether controlled OS alone provides fertility benefit for women with endometriosis and whether gonadotrophins provide benefit over, for example, letrozole (Aygen *et al.*, 2010).

Endometriosis may have a negative impact on IVF success rates compared with other causes of infertility (Barnhart *et al.*, 2002). Nonetheless, IVF is recommended as a fertility treatment for women with endometriosis, especially if fallopian tube function is compromised or if there are other infertility factors such as male factor (Soliman *et al.*, 1993). The chance of success is similar for GnRH antagonist versus GnRH agonist protocols (Benschop *et al.*, 2010). IVF does not appear to increase the risk of recurrence of endometriosis (D'Hooghe *et al.*, 2006).

Adjunct therapy to assisted conception for endometriosis-associated infertility

Medical treatment (including GnRH agonist) (Rickes *et al.*, 2002) and laparoscopic surgical treatment (Tanahatue *et al.*, 2005) prior to IUI/COS is not recommended, since there are insufficient data demonstrating benefit.

Treatment with GnRH agonist for 3–6 months prior to IVF is effective at improving the chance of IVF success (Sallam *et al.*, 2006). There are insufficient data to recommend the use of OCP prior to IVF/ICSI (de Ziegler *et al.*, 2010) and no data to compare the approach of pre-treatment with OCP versus GnRH agonist. There is concern that the presence of an endometrioma may damage the ovary, yet on the other hand, ovarian response to stimulation in IVF might be reduced in some women who have had an endometrioma removed (Yu *et al.*, 2010). The benefit of laparoscopic removal of endometriosis and/or endometriomas prior to IVF is unclear with respect to IVF outcome (Bianchi *et al.*, 2009; Benschop *et al.*, 2010), although it may improve access to the ovaries and even reduce the chance of infection related to the oocyte collection procedure. Whilst laparoscopic surgery following repeat failure of IVF treatment may improve the chance of natural conception, its role as an adjunct to IVF is unclear. Any decisions to perform surgery for endometriomas or deep endometriosis before ART should be made only after fully informed consent by surgeons with appropriate expertise.

Medical therapy for endometriosis-associated infertility

There is no evidence of fertility benefit from medical treatment; ovulation suppression may delay pregnancy and this is not recommended (Hughes *et al.*, 2007).

Emerging therapies for endometriosis-associated infertility

In one RCT, uterine bathing and tubal flushing with the oil-soluble contrast medium lipiodol has been reported to improve live birth rates in women, with endometriosis with otherwise unexplained infertility, who are attempting natural conception (Johnson *et al.*, 2004). The role of lipiodol hysterosalpingography as an adjunct to IVF remains unclear (Reilly *et al.*, 2011).

There is insufficient evidence to recommend the use of the following for fertility benefit: pentoxifylline (Lu *et al.*, 2012), traditional Chinese medicine (Guo *et al.*, 2010; Flower *et al.*, 2012), vitamins C or E (Mier-Cabrera *et al.*, 2008), mifepristone (Guo *et al.*, 2011), rosiglitazone (Moravek *et al.*, 2009) or valproic acid (Liu and Guo, 2008).

Discussion

We have developed a first international consensus statement on the management of endometriosis through rigorous methodology. An obvious finding in the quest for a consensus statement is that unanimity from a range of experts in any statement is difficult to attain. In our survey that followed the consensus meeting, none of the statements made achieved 100% agreement without the expression of a caveat about either the statement or the strength of the statement, and only 7 of our 69 consensus statements were associated with a 0% disagreement rate from the survey respondents. However, in the case of only two statements, we were unable to achieve a majority consensus.

The strength of this consensus statement is that it is truly international, with a breadth of representation from six continents across medical, surgical and fertility organizations, including a voice for the women themselves via 16 involved endometriosis organizations. There are potential weaknesses in a consensus process such as this. Some of our statements are not strongly based on research evidence and were termed GPPs; however, such statements could still be associated with a strong consensus amongst the group of experts. We will inevitably have overlooked some interventions that could be relevant, in spite of the methodology and feedback from all participants. It is therefore intended that this consensus will be updated regularly in response to feedback and, hopefully, increasing evidence in our field.

Unsurprisingly, there are similarities in our consensus statements with existing guidelines for managing endometriosis, but also the kind of differences that might be expected from the coalescence of an eclectic group of experts from many different standpoints. One of the real values to the participants in such an exercise is the opportunity to recognize a completely new perspective and interpretation of existing evidence; this can be applied in any multidisciplinary setting, where specialists in medical, surgical and fertility treatment join forces with women affected by endometriosis. In some cases, the strength of our statements (and in some cases, even the GRADE score) or the content of statements themselves conflict with those in other guidelines. We endeavoured to make strong statements (i) where the evidence was moderate or strong, in other words derived from reliable and reproducible RCTs (and even in some cases where the evidence was insufficient or negative where such evidence was deemed strong) or (ii) where the risk or expense of an intervention strongly justifies its non-use in the context of marginal or insufficient evidence or (iii) where there was enormous potential for benefit from a simple, low-invasive, low-cost intervention, to overcome a substantial burden of suffering, even in the face of only weak or absent research evidence (as in the case of our GPPs).

It must be emphasized that our process differed from that of guideline development. There is no general consensus on the most appropriate methodology for consensus statements and so we have adopted the methodology for the GRADE system of grading the quality of evidence (Guyatt *et al.*, 2008) (now felt to be the most relevant method of grading evidence and recommendations in guidelines) and adapted this to our consensus process. The turbulence that is present in the normal clinical environment is reflected by the fact that there is much lack of consensus amongst experts surrounding all aspects of the management of endometriosis. This also reflects the fact that the reality of the clinical situation at an individual level is far more complex than the idealized situation in an RCT. It must also be acknowledged that a consensus statement from international experts would almost certainly be subtly different with a different group of experts, although it is hoped that our broad sample of individuals was representative of the spectrum of viewpoints of all the members of all the organizations and societies represented.

Key issues that we have few answers for are management of the adolescent who has, or might have, endometriosis (more research is required and focus needs to be applied to management algorithms for young women and adolescents) as well as intervention strategies in the younger age group designed to prevent endometriosis; lifestyle and dietary interventions (where research evidence is largely absent); standardization of long-term strategies for prevention of recurrent endometriosis; clarification of management strategies, both surgical and medical, for women with deep endometriosis; development of standards of experience and expertise required for surgeons undertaking advanced laparoscopic endometriosis surgery; standardization of centres/networks of expertise with regard to definition, accreditation and longevity; development of models of care in low-resource settings and understanding endometriosis and its potential treatment after menopause. We have not addressed the important issue of diagnosis and classification of endometriosis, which would benefit from a similar international consensus approach. Individualization of every woman's care is an important factor in long-term management. Furthermore, it is possible that a subpopulation of women with endometriosis (depending on age, impact of symptoms, severity of disease, current or future fertility wishes, lifestyle factors, previous treatments and possibly disease markers) will benefit from some form of medical treatment to alter the course of this condition longer term (Vercellini *et al.*, 2011); the challenge is to identify these subpopulations and long-term management strategies. Further assessment of emerging therapies is also a key factor and this has been much neglected in recent times. It is of concern that, although many pre-clinical studies have shown positive results, very few have progressed to become phase II/III clinical trials, let alone proved to be effective (Guo *et al.*, 2009). In 2009, of 15 registered clinical trials in endometriosis, listed as completed, only three had been published, whilst the remaining 12 (80%) were unpublished (Guo *et al.*, 2009). More systematic and coordinated research effort and funding is required at an international level, so that any breakthrough treatment does not remain elusive, nor any research effort is ignored in order for others to continue to build upon results, be these positive or negative.

Conclusion

This consensus initiative, undertaken on a global scale through global collaboration by the WES, kept uppermost the goal of improving the quality

of life for women with endometriosis. This paper is the outcome of the first ever attempt to bring a global collaborative consensus to the management of endometriosis, reflecting the best scientific evidence available.

Supplementary data

Supplementary data are available at <http://humrep.oxfordjournals.org/>.

Acknowledgements

This consensus statement is dedicated to our friend and colleague Professor David Healy who contributed to and presented at the consensus meeting in Montpellier, but whose untimely passing in May 2012 has meant the loss of a visionary leader in our field.

Authors' roles

N.P.J. and L.H. accept full authorship responsibility on behalf of the WES Montpellier Consortium.

Funding

This workshop was organized by the WES. Financial support for attendees was provided by the organizations that they represented and these are listed in Table 1.

Conflict of interest

The views expressed in this consensus document represent the consensus of the individuals in the World Endometriosis Society Montpellier Consortium and not necessarily those of the organizations. The following disclosures were provided by participants: Mauricio Abrao: consultancy for Bayer Pharma; David Adamson: research grants from EMD Merck Serono, IBSA, LabCorp, Schering Plough, Auxogyn and shareholder in Advanced Reproductive Care; Catherine Allaire: consultant to and speaker for Bayer Pharma AG and speaker for Covidien; Christian Becker: research grant from Bayer Pharma AG; Deborah Bush: travel support from Bayer Pharma AG; Kristof Chwalisz: employed by Abbott Laboratories; Hilary Critchley: support for consultancy and research collaboration from Bayer Pharma AG, and support for consultancy from Preglem and Abbott Laboratories; Thomas D'Hooghe: consultancy/grants from Bayer, Merck, Merck Serono, Astellas, Arresto, Roche, Proteomika, Teva; Gerard Dunselman: advisor to Abbott Laboratories; Johannes Evers: department receives two unrestricted research grants from MSD Netherlands BV and Ferring Netherlands BV for research unrelated to the topic of this manuscript and Editor-in-Chief of Human Reproduction; Thomas Faustmann: employed by Bayer Pharma AG; Axel Forman: consultancy/travel grants from Bayer Pharma AG; Ian Fraser: consultancies and/or lectures, advisory board participation and research studies (around endometriosis, menstrual disorders and/or contraception) for Bayer Healthcare, Merck/MSD, Daiichi Sankyo and Vifor Pharma, with relevant funding for expenses and honoraria; Linda Giudice: academic associate for Quest Diagnostics; stock in Merck and Pfizer; Stephan Gordts: consultant for Karl Storz; Bernard Hedon: research grants from Ferring Pharmaceuticals and

Genevri; Lone Hummelshoj: consultancy/travel expenses from Bayer Pharma AG, Abbott Laboratories, Ferring Pharmaceuticals; Neil Johnson: consultancy/travel expenses from Bayer Pharma, Merck-Serono, and MSD; Ludwig Kiesel: consultant to Bayer Pharma AG, Abbott Laboratories, and Pregel; Alan Lam: director of the Centre for Advanced Reproductive Endosurgery, which is supported by Ethicon-Endosurgery, Stryker, Aesculap; Charles Miller: research grants from Covidien, Femasys, Olympus, Novartis, Abbott, consultant to Ferring Pharmaceuticals, Ethicon Endo-Surgery, Ethicon Women's Health and Urology, Boston Scientific, Care Fusion, Hologic, Covidien, Femasys, Abbott, Halt Medical, speaker for Ferring Pharmaceuticals, Ethicon Endo-Surgery, Ethicon Women's Health and Urology, Merck, Femasys, Smith & Nephew; Carlos Petta: consultant to Bayer Pharma AG, Merck, and MSD; Edgardo Rolla: travel expenses from Bayer Pharma AG, Laboratorios Bagó SA, and MSD; Luk Rombauts: educational grants, consultancy fees and unrestricted research funding from Merck-Serono and MSD and minor shareholder of Monash IVF; Sony Singh: speaker/consultant for Bayer Pharma and Abbott Industries; David Soriano: consultancy Bayer Pharma; Jim Tsaltas: travel and research grants from Merck Serono; fellowship funding from Covidien. The following disclosed no competing interests: Elisabet Andersson, Kolbrún Birna Árdal, Bianca de Bie, Cindy Farquhar, Heather Guidone, Sun-Wei Guo, Johanna Hulkkonen, Louise Hull, Miriam Just, Clodagh Lynam, Lise-lotte Mettler, Helen North, Lucy Prentice, Shelley Reilly, Fernando Reis, Karl-Werner Schweppe, Kathy Sharpe-Timms, Dian Shepperson Mills, Martyn Stafford-Bell, Pamela Stratton, Robert Taylor. The following made no declaration: Vibeke Amelung, Mary-Lou Ballweg, Jessica Fourquet, David Healy, Rishma Pai, Tamer Seckin.

References

- Abbott J, Hawe J, Hunter D, Holmes M, Finn P, Garry R. Laparoscopic excision of endometriosis: a randomized, placebo-controlled trial. *Fertil Steril* 2004;**82**:878–884.
- Abou-Setta AM, Al-Inany HG, Farquhar CM. Levonorgestrel-releasing intrauterine device (LNG-IUD) for symptomatic endometriosis following surgery. *Cochrane Database Syst Rev* 2006;**4**:CD005072.
- Adamson GD. Endometriosis classification: an update. *Curr Opin Obstet Gynecol* 2011;**23**:213–220.
- Adamson GD, Pasta DJ. Endometriosis fertility index: the new, validated endometriosis staging system. *Fertil Steril* 2010;**94**:1609–1615.
- Adamson GD, Kennedy SH, Hummelshoj L. Creating solutions in endometriosis: global collaboration through the World Endometriosis Research Foundation. *J Endometriosis* 2010;**2**:3–6.
- Al-Jefout M, Dezarnaulds G, Cooper M, Tokushige N, Luscombe GM, Markham R, Fraser IS. Diagnosis of endometriosis by detection of nerve fibres in an endometrial biopsy: a double blind study. *Hum Reprod* 2009;**24**:3019–3024.
- Al Kadri H, Hassan S, Al-Fozan HM, Hajeer A. Hormone therapy for endometriosis and surgical menopause. *Cochrane Database Syst Rev* 2009;**1**:CD005997.
- Allen C, Hopewell S, Prentice A, Gregory D. Nonsteroidal anti-inflammatory drugs for pain in women with endometriosis. *Cochrane Database Syst Rev* 2009;**2**:CD004753.
- Aygen MA, Atakul T, Oner G, Ozgun MT, Sahin Y, Ozturk F. A prospective trial comparing letrozole versus gonadotropins for ovulation induction in patients with histologically documented endometriosis. *Hum Reprod* 2010;**25**(S1):6.
- Barnhart K, Dunsmoor-Su R, Coutifaris C. Effect of endometriosis on in vitro fertilization. *Fertil Steril* 2002;**77**:1148–1155.
- Barri PN, Coroleu B, Tur R, Barri-Soldevila PN, Rodríguez I. Endometriosis-associated infertility: surgery and IVF, a comprehensive therapeutic approach. *Reprod Biomed Online* 2010;**21**:179–185.
- Bayoglu TY, Dilbaz B, Altinbas SK, Dilbaz S. Postoperative medical treatment of chronic pelvic pain related to severe endometriosis: levonorgestrel-releasing intrauterine system versus gonadotropin-releasing hormone analogue. *Fertil Steril* 2011;**95**:492–496.
- Benschop L, Farquhar C, Van der Poel N, Heineman MJ. Interventions for women with endometrioma prior to assisted reproductive technology. *Cochrane Database Syst Rev* 2010;**11**:CD008571.
- Biacchiardi CP, Plane LD, Camanni M, Deltetto F, Delpiano EM, Marchino GL, Gennarelli G, Revelli A. Laparoscopic stripping of endometriomas negatively affects ovarian follicular reserve even if performed by experienced surgeons. *Reprod Biomed Online* 2011;**23**:740–746.
- Bianchi PH, Pereira RM, Zanatta A, Alegretti JR, Motta EL, Serafini PC. Extensive excision of deep infiltrative endometriosis before in vitro fertilization significantly improves pregnancy rates. *J Minim Invasive Gynecol* 2009;**16**:174–180.
- Bokor A, Kyama CM, Vercruyse L, Fassbender A, Gevaert O, Vodolazkaia A, De Moor B, Fülöp V, D'Hooghe T. Density of small diameter sensory nerve fibres in endometrium: a semi-invasive diagnostic test for minimal to mild endometriosis. *Hum Reprod* 2009;**24**:3025–3032.
- Brosens I, Brosens JJ, Fusi L, Al-Sabbagh M, Kuroda K, Benagiano G. Risks of adverse pregnancy outcome in endometriosis. *Fertil Steril* 2012;**98**:30–35.
- Brown J, Pan A, Hart RJ. Gonadotrophin-releasing hormone analogues for pain associated with endometriosis. *Cochrane Database Syst Rev* 2010;**12**:CD008475.
- Brown J, Kives S, Akhtar M. Progestagens and anti-progestagens for pain associated with endometriosis. *Cochrane Database Syst Rev* 2012;**3**:CD002122.
- Bush D. Endometriosis New Zealand—a national organisation pivotal to facilitating the wellness of women. In: Rombauts L (ed.). *World Endometriosis Soc eJournal* 2009;**11**:6–9.
- Chapron C, Fritel X, Dubuisson JB. Fertility after laparoscopic management of deep endometriosis infiltrating the uterosacral ligaments. *Hum Reprod* 1999;**14**:329–332.
- Cosson M, Querleu D, Donnez J, Madelenat P, Koninckx P, Audebert A, Manhes H. Dienogest is as effective as triptorelin in the treatment of endometriosis after laparoscopic surgery: results of a prospective, multicenter, randomized study. *Fertil Steril* 2002;**77**:684–692.
- Costello MF. Systematic review of the treatment of ovulatory infertility with clomiphene citrate and intrauterine insemination. *Aust N Z J Obstet Gynaecol* 2004;**44**:93–102.
- Crosignani PG, Luciano A, Ray A, Bergqvist A. Subcutaneous depot medroxyprogesterone acetate versus leuprolide acetate in the treatment of endometriosis-associated pain. *Hum Reprod* 2006;**21**:248–256.
- Davis L, Kennedy SS, Moore J, Prentice A. Modern combined oral contraceptives for pain associated with endometriosis. *Cochrane Database Syst Rev* 2007;**3**:CD001019.
- de Ziegler D, Gayet V, Aubriot FX, Fauque P, Streuli I, Wolf JP, de Mouzon J, Chapron C. Use of oral contraceptives in women with endometriosis before assisted reproduction treatment improves outcomes. *Fertil Steril* 2010;**94**:2796–2799.
- D'Hooghe T, Hummelshoj L. Multi-disciplinary centres/networks of excellence for endometriosis management and research: a proposal. *Hum Reprod* 2006;**21**:2743–2748.

- D'Hooghe TM, Denys B, Spiessens C, Meuleman C, Debrock S. Is the endometriosis recurrence rate increased after ovarian hyperstimulation? *Fertil Steril* 2006;**86**:283–290.
- Donnez J, Squifflet J. Complications, pregnancy and recurrence in a prospective series of 500 patients operated on by the shaving technique for deep rectovaginal endometriotic nodules. *Hum Reprod* 2010;**25**:1949–1958.
- Dovey S, Sanfilippo J. Endometriosis and the adolescent. *Clin Obstet Gynecol* 2010;**53**:420–428.
- Fagervold B, Jenssen M, Hummelshoj L, Moen MH. Life after a diagnosis with endometriosis - a 15 years follow-up study. *Acta Obstet Gynecol Scand* 2009;**88**:914–919.
- Farmer JE, Prentice A, Breeze A, Ahmad G, Duffy JMN, Watson A, Pick A. Gonadotrophin-releasing hormone analogues for endometriosis: bone mineral density. *Cochrane Database Syst Rev* 2009;**1**:CD001297.
- Fernando S, Breheny S, Jaques AM, Halliday JL, Baker G, Healy D. Preterm birth, ovarian endometriomata, and assisted reproduction technologies. *Fertil Steril* 2009;**91**:325–330.
- Ferrero S, Anserini P, Abbamonte LH, Ragni N, Camerini G, Remorgida V. Fertility after bowel resection for endometriosis. *Fertil Steril* 2009;**92**:41–46.
- Ferrero S, Gillott DJ, Venturini PL, Remorgida V. Use of aromatase inhibitors to treat endometriosis-related pain symptoms: a systematic review. *Reprod Biol Endocrinol* 2011;**9**:89.
- Flower A, Liu JP, Lewith G, Little P, Li Q. Chinese herbal medicine for endometriosis. *Cochrane Database Syst Rev* 2012;**5**:CD006568.
- Ford J, English J, Miles WA, Giannopolous T. Pain, quality of life and complications following the radical resection of rectovaginal endometriosis. *Br J Obstet Gynaecol* 2004;**111**:353–356.
- French SD, Cameron M, Walker BF, Reggars JW, Esterman AJ. Superficial heat or cold for low back pain. *Cochrane Database of Syst Rev* 2006;**1**:CD004750.
- Furness S, Yap C, Farquhar C, Cheong YC. Pre and post-operative medical therapy for endometriosis surgery. *Cochrane Database Syst Rev* 2009;**1**:CD003678.
- Giudice LC. Clinical practice: endometriosis. *N Engl J Med* 2010;**362**:2389–2398.
- Guo SW. Recurrence of endometriosis and its control. *Hum Reprod Update* 2009;**15**:441–461.
- Guo SW, Hummelshoj L, Olive DL, Bulun SE, D'Hooghe TM, Evers JL. A call for more transparency of registered clinical trials on endometriosis. *Hum Reprod* 2009;**24**:1247–1254.
- Guo SW, He W, Zhao T, Liu X, Zhang T. Clinical trials and trial-like studies on the use of traditional Chinese medicine to treat endometriosis. *Expert Rev Obstet Gynecol* 2010;**5**:533–555.
- Guo SW, Liu M, Shen F, Liu X. Use of mifepristone to treat endometriosis: a review of clinical trials and trial-like studies conducted in China. *Womens Health* 2011;**7**:51–70.
- Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, Schünemann HJ, GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;**336**:924–926.
- Guzick DS, Huang LS, Broadman BA, Nealon M, Hornstein MD. Randomized trial of leuprolide versus continuous oral contraceptives in the treatment of endometriosis-associated pelvic pain. *Fertil Steril* 2011;**95**:1568–1573.
- Harada T, Momoeda M, Taketani Y, Hoshiai H, Terakawa N. Low-dose oral contraceptive pill for dysmenorrhea associated with endometriosis: a placebo-controlled, double-blind, randomized trial. *Fertil Steril* 2008;**90**:1583–1588.
- Harada T, Momoeda M, Taketani Y, Aso T, Fukunaga M, Hagino H, Terakawa N. Dienogest is as effective as intranasal buserelin acetate for the relief of pain symptoms associated with endometriosis—a randomized, double-blind, multicenter, controlled trial. *Fertil Steril* 2009;**91**:675–681.
- Hart RJ, Hickey M, Maouris P, Buckett W. Excisional surgery versus ablative surgery for ovarian endometriomata. *Cochrane Database Syst Rev* 2008;**2**:CD004992.
- Healey M, Ang WC, Cheng C. Surgical treatment of endometriosis: a prospective randomized double-blinded trial comparing excision and ablation. *Fertil Steril* 2010;**94**:2536–2540.
- Hughes E, Brown J, Collins JJ, Farquhar C, Fedorkow DM, Vanderkerchove P. Ovulation suppression for endometriosis for women with subfertility. *Cochrane Database Syst Rev* 2007;**3**:CD000155.
- Jacobson TZ, Duffy JM, Barlow D, Koninckx PR, Garry R. Laparoscopic surgery for pelvic pain associated with endometriosis. *Cochrane Database Syst Rev* 2009;**4**:CD001300.
- Jacobson TZ, Duffy JM, Barlow D, Farquhar C, Koninckx PR, Olive D. Laparoscopic surgery for subfertility associated with endometriosis. *Cochrane Database Syst Rev* 2010;**1**:CD001398.
- Johnson NP, Farquhar CM, Hadden WE, Suckling J, Yu Y, Sadler L. The FLUSH trial—flushing with lipiodol for unexplained (and endometriosis-related) subfertility by hysterosalpingography: a randomized trial. *Hum Reprod* 2004;**19**:2043–2051.
- Kennedy S, Bergqvist A, Chapron C, D'Hooghe T, Dunselman G, Greb R, Hummelshoj L, Prentice A, Saridogan E, ESHRE Special Interest Group for Endometriosis and Endometrium Guideline Development Group. ESHRE guideline for the diagnosis and treatment of endometriosis. *Hum Reprod* 2005;**20**:2698–2704.
- Köhler G, Faustmann TA, Gerlinger C, Seitz C, Mueck AO. A dose-ranging study to determine the efficacy and safety of 1, 2, and 4mg of dienogest daily for endometriosis. *Int J Gynaecol Obstet* 2010;**108**:21–25.
- Koninckx PR, Timmermans B, Meuleman C, Penninckx F. Complications of CO₂-laser endoscopic excision of deep endometriosis. *Hum Reprod* 1996;**11**:2263–2268.
- Koninckx PR, Craessaerts M, Timmerman D, Cornillie F, Kennedy S. Anti-TNF-alpha treatment for deep endometriosis-associated pain: a randomized placebo-controlled trial. *Hum Reprod* 2008;**23**:2017–2023.
- Koninckx PR, Ussia A, Adamyan L, Wattiez A, Donnez J. Deep endometriosis: definition, diagnosis, and treatment. *Fertil Steril* 2012;**98**:564–571.
- Koppan A, Hamori J, Vranics I, Garai J, Kriszbacher I, Bodis J, Rebek-Nagy G, Koppan M. Pelvic pain in endometriosis: painkillers or sport to alleviate symptoms? *Acta Physiol Hung* 2010;**97**:234–239.
- Kroon B, Johnson N, Chapman M, Yazdani A, Hart R, Australasian CREI Consensus Expert Panel on Trial evidence (ACCEPT) group. Fibroids in infertility—consensus statement from ACCEPT (Australasian CREI Consensus Expert Panel on Trial evidence). *Aust N Z J Obstet Gynaecol* 2011;**51**:289–295.
- Laschke MW, Menger MD. Anti-angiogenic treatment strategies for the therapy of endometriosis. *Hum Reprod Update* 2012;**18**:682–702.
- Lauffer MR, Goitein L, Bush M, Cramer DW, Emans SJ. Prevalence of endometriosis in adolescent girls with chronic pelvic pain not responding to conventional therapy. *J Pediatr Adolesc Gynecol* 1997;**10**:199–202.
- Lessey DA. Assessment of endometrial receptivity. *Fertil Steril* 2011;**96**:522–529.
- Liu X, Guo SW. A pilot study on the off-label use of valproic acid to treat adenomyosis. *Fertil Steril* 2008;**89**:246–250.
- Lu D, Song H, Li Y, Clarke J, Shi G. Pentoxifylline for endometriosis. *Cochrane Database Syst Rev* 2012;**1**:CD007677.
- Lucas P. Cannabis as an adjunct to or substitute for opiates in the treatment of chronic pain. *J Psychoactive Drugs* 2012;**44**:125–133.

- Maubon A, Faury A, Kapella M, Pouquet M, Piver P. Uterine junctional zone at magnetic resonance imaging: a predictor of in vitro fertilization implantation failure. *J Obstet Gynaecol Res* 2010;**36**:611–618.
- May KE, Conduit-Hulbert SA, Villar J, Kirtley S, Kennedy SH, Becker CM. Peripheral biomarkers of endometriosis: a systematic review. *Hum Reprod Update* 2010;**16**:651–674.
- Mier-Cabrera J, Genera-Garcia M, De la Jara-Diaz J, Perichart-Perera O, Vellido-Ortega F, Hernandez-Guerrero C. Effect of vitamins C and E supplementation on peripheral oxidative stress markers and pregnancy rate in women with endometriosis. *Int J Gynaecol Obstet* 2008;**100**:252–256.
- Moen MH, Rees M, Brincat M, Erel T, Gambacciani M, Lambrinoudaki I, Schenck-Gustafsson K, Tremollieres F, Vujovic S, Rozenberg S, European Menopause and Andropause Society. EMAS position statement: Managing the menopause in women with a past history of endometriosis. *Maturitas* 2010;**67**:94–97.
- Momoeda M, Harada T, Terakawa N, Aso T, Fukunaga M, Hagino H, Taketani Y. Long-term use of dienogest for the treatment of endometriosis. *J Obstet Gynaecol Res* 2009;**35**:1069–1076.
- Moravek MB, Ward EA, Lebovic DI. Thiazolidinediones as therapy for endometriosis: a case series. *Gynecol Obstet Invest* 2009;**68**:167–170.
- Mutihir JT, Nyango DD. Massive haemoperitoneum from endometriosis masquerading as ruptured ectopic pregnancy: case report. *Niger J Clin Pract* 2010;**13**:477–479.
- Nnoaham K, Hummelshoj L, Webster P, D'Hooghe T, de Cicco Nardone F, de Cicco Nardone C, Jenkinson C, Kennedy SH, Zondervan KT, World Endometriosis Research Foundation Global Study of Women's Health Consortium. Impact of endometriosis on quality of life and work productivity: A multi-centre study across 10 countries. *Fertil Steril* 2011;**96**:366–373.
- Nnoaham KE, Hummelshoj L, Kennedy SH, Jenkinson C, Zondervan KT, World Endometriosis Research Foundation Women's Health Symptom Survey Consortium. Developing symptom-based predictive models of endometriosis as a clinical screening tool: results from a multicenter study. *Fertil Steril* 2012;**98**:692–701.
- Pabona JM, Simmen FA, Nikiforov MA, Zhuang D, Shankar K, Velarde MC, Zelenko Z, Giudice LC, Simmen RC. Krüppel-like factor 9 and progesterone receptor coregulation of decidualizing endometrial stromal cells: implications for the pathogenesis of endometriosis. *J Clin Endocrinol Metab* 2012;**97**:E376–E392.
- Pearce CL, Templeman C, Rossing MA, Lee A, Near AM, Webb PM, Nagle CM, Doherty JA, Cushing-Haugen KL, Wicklund KG et al. Association between endometriosis and risk of histological subtypes of ovarian cancer: a pooled analysis of case-control studies. *Lancet Oncol* 2012;**13**:385–394.
- Pellicano M, Bramante S, Guida M, Bifulco G, Di Spiezio Sardo A, Cirillo D, Nappi C. Ovarian endometrioma: postoperative adhesions following bipolar coagulation and suture. *Fertil Steril* 2008;**89**:796–799.
- Petraglia F, Hornung D, Seitz C, Faustmann T, Gerlinger C, Luisi S, Lazzeri L, Strowitzki T. Reduced pelvic pain in women with endometriosis: efficacy of long-term dienogest treatment. *Arch Gynecol Obstet* 2012;**285**:167–173.
- Proctor M, Murphy PA. Herbal and dietary therapies for primary and secondary dysmenorrhoea. *Cochrane Database Syst Rev* 2001;**2**:CD002124.
- Proctor ML, Smith CA, Farquhar CM, Stones RW. Transcutaneous electrical nerve stimulation and acupuncture for primary dysmenorrhoea. *Cochrane Database Syst Rev* 2002;**1**:CD002123.
- Proctor ML, Latthe PM, Farquhar CM, Khan KS, Johnson NP. Surgical interruption of pelvic nerve pathways for primary and secondary dysmenorrhoea. *Cochrane Database Syst Rev* 2005;**4**:CD001896.
- Proctor M, Hing W, Johnson TC, Murphy PA, Brown J. Spinal manipulation for dysmenorrhoea. *Cochrane Database Syst Rev* 2006;**3**:CD002119.
- Proctor M, Murphy PA, Pattison HM, Suckling JA, Farquhar C. Behavioural interventions for dysmenorrhoea. *Cochrane Database Syst Rev* 2007;**3**:CD002248.
- Reilly SJ, Stewart AW, Prentice LR, Johnson NP. The IVF-LUBE trial: lipidol uterine bathing effect for enhancing the results of in vitro fertilisation, a pilot randomised trial. In: *11th World Congress on Endometriosis*, 2011;p27:S#10–5.
- Revised American Society for Reproductive Medicine. Revised American Society for Reproductive Medicine classification of endometriosis: 1996. *Fertil Steril* 1997;**67**:817–821.
- Rickes D, Nickel I, Kropf S, Kleinstein J. Increased pregnancy rates after ultralong postoperative therapy with gonadotropin-releasing hormone analogs in patients with endometriosis. *Fertil Steril* 2002;**78**:757–762.
- Roman JD. Adolescent endometriosis in the Waikato region of New Zealand – a comparative cohort study with a mean follow up time of 2.6 years. *Aust N Z J Obstet Gynaecol* 2010;**50**:579–583.
- Sallam HN, Garcia-Velasco JA, Dias S, Arici A. Long-term pituitary down-regulation before in vitro fertilization (IVF) for women with endometriosis. *Cochrane Database Syst Rev* 2006;**1**:CD004635.
- Sayasneh A, Tsivos D, Crawford R. Endometriosis and ovarian cancer: a systematic review. *Obstet Gynecol* 2011;**2011**:1403–1410.
- Schlaff WD, Carson SA, Luciano A, Ross D, Bergqvist A. Subcutaneous injection of depot medroxyprogesterone acetate compared with leuprolide acetate in the treatment of endometriosis-associated pain. *Fertil Steril* 2006;**85**:314–325.
- Schweppe KW, Ring D. Peritoneal defects and the development of endometriosis in relation to the timing of endoscopic surgery during the menstrual cycle. *Fertil Steril* 2002;**78**:763–766.
- Selak V, Farquhar C, Prentice A, Singla A. Danazol for pelvic pain associated with endometriosis. *Cochrane Database Syst Rev* 2007;**4**:CD000068.
- Seracchioli R, Mabrouk M, Frascà C, Manuzzi L, Montanari G, Keramyda A, Venturoli S. Long-term cyclic and continuous oral contraceptive therapy and endometrioma recurrence: a randomized controlled trial. *Fertil Steril* 2010;**93**:52–56.
- Sesti F, Pietropolli A, Capozzolo T, Broccoli P, Pierangeli S, Bollea MR, Piccione E. Hormonal suppression treatment or dietary therapy versus placebo in the control of painful symptoms after conservative surgery for endometriosis stage III-IV. A randomized comparative trial. *Fertil Steril* 2007;**88**:1541–1547.
- Sesti F, Capozzolo T, Pietropolli A, Marziali M, Bollea MR, Piccione E. Recurrence rate of endometrioma after laparoscopic cystectomy: a comparative randomized trial between post-operative hormonal suppression treatment or dietary therapy vs. placebo. *Eur J Obstet Gynecol Reprod Biol* 2009;**147**:72–77.
- Shakiba K, Bena JF, McGill KM, Minger J, Falcone T. Surgical treatment of endometriosis: a 7-year follow-up on the requirement for further surgery. *Obstet Gynecol* 2008;**111**:1285–1292.
- Simoens S, Hummelshoj L, D'Hooghe T. Endometriosis: cost estimates and methodological perspective. *Hum Reprod Update* 2007;**13**:395–404.
- Simoens S, Dunselman G, Dirksen C, Hummelshoj L, Bokor A, Brandes I, Brodsky V, Canis M, Colombo GL, DeLeire T et al. The burden of endometriosis: costs and quality of life of women with endometriosis and treated in referral centres. *Hum Reprod* 2012;**27**:1292–1299.
- Soliman S, Daya S, Collins J, Jarrell J. A randomized trial of in vitro fertilization versus conventional treatment for infertility. *Fertil Steril* 1993;**59**:1239–1244.
- Somigliana E, Berlanda N, Benaglia L, Viganò P, Vercellini P, Fedele L. Surgical excision of endometriomas and ovarian reserve: a systematic

- review on serum antimüllerian hormone level modifications. *Fertil Steril* 2012;**98**:1531–1538.
- Spitz IM. Clinical utility of progesterone receptor modulators and their effect on the endometrium. *Curr Opin Obstet Gynecol* 2009;**21**: 318–324.
- Stephansson O, Kieler H, Granath F, Falconer H. Endometriosis, assisted reproduction technology and risk of adverse pregnancy outcome. *Hum Reprod* 2009;**24**:2341–2347.
- Stepniowska A, Pomini P, Scioscia M, Mereu L, Ruffo G, Minelli L. Fertility and clinical outcome after bowel resection in infertile women with endometriosis. *Reprod Biomed Online* 2010;**20**:602–609.
- Stratton P, Berkley KJ. Chronic pelvic pain and endometriosis: translational evidence of the relationship and implications. *Hum Reprod Update* 2011; **17**:327–346.
- Stratton P, Sinaii N, Segars J, Koziol D, Wesley R, Zimmer C, Winkel C, Nieman LK. Return of chronic pelvic pain from endometriosis after raloxifene treatment: a randomized controlled trial. *Obstet Gynecol* 2008;**111**:88–96.
- Streuli I, de Ziegler D, Gayet V, Santulli P, Bijaoui G, de Mouzon J, Chapron C. In women with endometriosis anti-Müllerian hormone levels are decreased only in those with previous endometrioma surgery. *Hum Reprod* 2012;**27**:3294–3303.
- Strowitzki T, Faustmann T, Gerlinger C, Seitz C. Dienogest in the treatment of endometriosis-associated pelvic pain: a 12-week, randomized, double-blind, placebo-controlled study. *Eur J Obstet Gynecol Reprod Biol* 2010a;**151**:193–198.
- Strowitzki T, Marr J, Gerlinger C, Faustmann T, Seitz C. Dienogest is as effective as leuprolide acetate in treating the painful symptoms of endometriosis: a 24-week, randomized, multicentre, open-label trial. *Hum Reprod* 2010b;**25**:633–641.
- Strowitzki T, Marr J, Gerlinger C, Faustmann T, Seitz C. Detailed analysis of a randomized, multicenter, comparative trial of dienogest versus leuprolide acetate in endometriosis. *Int J Gynaecol Obstet* 2012; **117**:228–233.
- Struthers RS, Nicholls AJ, Grundy J, Chen T, Jimenez R, Yen SS, Bozigian HP. Suppression of gonadotropins and estradiol in premenopausal women by oral administration of the nonpeptide gonadotropin-releasing hormone antagonist elagolix. *J Clin Endocrinol Metab* 2009;**94**:545–551.
- Subit M, Gantt P, Broce M, Seybold D, Randall D. Endometriosis associated infertility: double intrauterine insemination improves fecundity in patients positive for antiendometrial antibodies. *Am J Reprod Immunol* 2011;**66**:100–107.
- Tanahatoo SJ, Lambalk CB, Hompes PG. The role of laparoscopy in intrauterine insemination: a prospective randomized reallocation study. *Hum Reprod* 2005;**20**:3225–3230.
- Tsolakidis D, Pados G, Vavilis D, Athanatos D, Tsalikis T, Giannakou A, Tarlatzis BC. The impact on ovarian reserve after laparoscopic ovarian cystectomy versus three-stage management in patients with endometriomas: a prospective randomized study. *Fertil Steril* 2010; **94**:71–77.
- Tummon I, Asher L, Martin J, Tulandi T. Randomized controlled trial of superovulation and insemination for infertility associated with minimal or mild endometriosis. *Fertil Steril* 1997;**68**:8–12.
- Vercellini P, Pietropaolo G, De Giorgi O, Daguati R, Pasin R, Crosignani PG. Reproductive performance in infertile women with rectovaginal endometriosis: is surgery worthwhile? *Am J Obstet Gynecol* 2006;**195**:1303–1310.
- Vercellini P, Somigliana E, Viganò P, De Matteis S, Barbara G, Fedele L. The effect of second-line surgery on reproductive performance of women with recurrent endometriosis: a systematic review. *Acta Obstet Gynecol Scand* 2009;**88**:1074–1082.
- Vercellini P, Giussy B, Somigliana E, Bianchi S, Abbiati A, Fedele L. Comparison of contraceptive ring and patch for the treatment of symptomatic endometriosis. *Fertil Steril* 2010;**93**:2150–2161.
- Vercellini P, Crosignani P, Somigliana E, Viganò P, Frattaruolo MP, Fedele L. Waiting for Godot: a commonsense approach to the medical treatment of endometriosis. *Hum Reprod* 2011;**26**:3–13.
- Vercellini P, De Matteis S, Somigliana E, Buggio L, Frattaruolo MP, Fedele L. Long-term adjuvant therapy for the prevention of postoperative endometrioma recurrence: a systematic review and meta-analysis. *Acta Obstet Gynecol Scand* 2013;**92**:8–16.
- Wright J, Lotfallah H, Jones K, Lovell D. A randomized trial of excision versus ablation for mild endometriosis. *Fertil Steril* 2005;**83**:1830–1836.
- Wykes CB, Clark TJ, Khan KS. Accuracy of laparoscopy in the diagnosis of endometriosis: a systematic quantitative review. *Brit J Obstet Gynaecol* 2004;**111**:1204–1212.
- Yeung P Jr, Sinervo K, Winer W, Albee RB Jr. Complete laparoscopic excision of endometriosis in teenagers: is postoperative hormonal suppression necessary? *Fertil Steril* 2011;**95**:1909–1912.
- Yu HT, Huang HY, Soong YK, Lee CL, Chao A, Wang CJ. Laparoscopic ovarian cystectomy of endometriomas: surgeons' experience may affect ovarian reserve and live-born rate in infertile patients with in vitro fertilization-intracytoplasmic sperm injection. *Eur J Obstet Gynecol Reprod Biol* 2010;**152**:172–175.
- Zhu X, Proctor M, Bensoussan A, Wu E, Smith CA. Chinese herbal medicine for primary dysmenorrhoea. *Cochrane Database Syst Rev* 2008;**2**:CD005288.
- Zhu X, Hamilton KD, McNicol ED. Acupuncture for pain in endometriosis. *Cochrane Database Syst Rev* 2011;**9**:CD007864.

Appendix

The complete list of people representing The World Endometriosis Society Montpellier Consortium is as follows: M.S. Abrao, G.D. Adamson, C. Allaire, V. Amelung, E. Andersson, C. Becker, K.B. Birna Árdal, D. Bush, B. de Bie, K. Chwalisz, H. Critchley, T. D'Hooghe, G. Dunselman, J.L.H. Evers, C. Farquhar, T. Faustmann, A. Forman, J. Fourquet, I. Fraser, L. Giudice, S. Gordts, H. Guidone, S.W. Guo, D. Healy, B. Hedon, J. Hulkkonen, L. Hull, L. Hummelshoj, N.P. Johnson, M. Just, L. Kiesel, A. Lam, C. Lynam, L. Mettler, C. Miller, H. North, R. Pai, C. Petta, L. Prentice, S. Reilly, F. Reis, E. Rolla, L. Rombauts, K.W. Schweppe, T. Seckin, K. Sharpe-Timms, D. Shepperson Mills, S. Singh, D. Soriano, M. Stafford-Bell, P. Stratton, R. Taylor, J. Tsaltas, J. Veit and P. Vercellini.